

BD

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<p>(51) International Patent Classification <sup>6</sup> : <b>C07K 14/62, 1/30</b></p>	<p><b>A1</b></p>	<p>(11) International Publication Number: <b>WO 98/34953</b> (43) International Publication Date: <b>13 August 1998 (13.08.98)</b></p>
<p>(21) International Application Number: <b>PCT/DK98/00042</b> (22) International Filing Date: <b>3 February 1998 (03.02.98)</b> (30) Priority Data: <b>0146/97 7 February 1997 (07.02.97) DK</b> (71) Applicant: <b>NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsvaerd (DK).</b> (72) Inventors: <b>WHITTINGHAM, Jean, Lesley; 16 Markham Street, Haxby Road, York YO3 7NR (GB). BALSCHMIDT, Per; Tibberup Allé 20, DK-3060 Espergærde (DK).</b></p>	<p>(81) Designated States: <b>AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GR, GH, GM, GW, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b>  Published <i>With international search report.</i></p>	
<p>(54) Title: <b>CRYSTALLISATION OF PROTEINS</b> (57) Abstract  A method of providing zinc containing crystals of a protein derivative which has a lysine residue which carries a lipophilic substituent on the <math>\epsilon</math>-amino group, said method comprising: a) providing a solution of the protein derivative in an alkaline buffer, which further contains a zinc salt; b) adjusting the pH value of the solution to a value between 7 and 10 and c) isolating the crystals formed.</p>		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SS	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## CRYSTALLISATION OF PROTEINS

## FIELD OF THE INVENTION

- 5 The present invention relates to a method of providing filterable crystals of zinc complexes of a protein which has a lysine residue which carries a lipophilic substituent on the  $\epsilon$ -amino group. In particular, the present invention relates to a method of providing crystals of zinc complexes of proinsulins, insulins and insulin analogues which have a lysine residue which carries a lipophilic substituent on the  $\epsilon$ -amino group. Optionally, the crystals also contain a  
10 phenol.

## BACKGROUND OF THE INVENTION

- 15 The isolation of pharmaceutical proteins in the crystalline state is important because crystals can be dried easily and subsequently stored at low temperature under conditions where the stability of the bulk protein is optimal. Insulins and insulin analogues in which the  $\epsilon$ -amino group of a lysine residue contained therein has a lipophilic substituent e.g. in the form of an acyl group have a protracted profile of action and show promise for use in long-acting basal  
20 therapy in the treatment of IDDM (insulin-demanding diabetes mellitus) and NIDDM (non-insulin-demanding diabetes mellitus). The preparation of such acylated insulins and insulin analogues is described *i.a.* in Japanese patent application 1-254,699 (Kodama), in WO 95/07931 (Novo Nordisk), in EP 0 712 862 A2 (Eli Lilly) and in WO 96/29344 (Novo Nordisk). Unfortunately, such acylated insulins and insulin analogues have been found to be less  
25 prone to crystallize than the unmodified parent compounds.

- Proinsulins, insulins and insulin analogues are rather labile proteins and their stability depends *i.a.* on the purity of the particular preparation. Optimal stability can be expected when a pure preparation is kept in solid form, in particular in the form of crystals. Unfortunately,  
30 precipitation of these compounds from solutions usually leads to amorphous precipitates which are difficult or impossible to isolate by filtration. Instead they can be isolated by centrifugation. However, when they are isolated by centrifugation it is difficult to free the particles efficiently from the mother liquor. Even when amorphous particles can be isolated by filtra-

tion, the amorphous material will usually be less pure than corresponding crystals because more impurities are embedded in amorphous material than in crystals.

The preparation of crystals of N<sup>629</sup>-(myristoyl)des(B30) human insulin is described in European patent application No. 94926816.3, Example 33. These zinc-free crystals were precipitated at pH 9 from 20% aqueous ethanol containing 0.625 M sodium chloride. A method for recovering acylated proteins, especially certain fatty acid-acylated insulins, by precipitation and filtration as a freely-flowing powder is described in EP 0 747 391 A2 (Eli Lilly). According to this method a filterable precipitate of a protein is obtained by adjusting the pH of an aqueous solution containing the protein and adding a suitable amount of alcohol. No formation of crystals is mentioned.

For use in therapy, proteins must be prepared in highly purified form and the storage conditions must ensure that degradation during the storage is minimized. An acylated insulin or an acylated insulin analogue in highly purified form is administered in the form of a solution of a zinc complex thereof in a composition which further comprises a phenolic compound. Accordingly, it would be convenient in the production to store the bulk acylated insulin or acylated insulin analogue in the form of filterable crystals containing both insulin or insulin analogue, zinc and a phenolic compound.

It is thus an object of the present invention to provide a method by which zinc complexes of an acylated insulin or an acylated insulin analogue, optionally also containing a phenolic compound, can be obtained in filterable, crystalline form.

According to the invention this object has been accomplished by precipitating the acylated insulin or acylated insulin analogue from an aqueous buffer.

#### SUMMARY OF THE INVENTION

Thus, in its broadest aspect, the present invention relates to a method of providing zinc containing crystals of a protein derivative which has a lysine residue which carries a lipophilic substituent on the  $\epsilon$ -amino group, said method comprising

- a) providing a solution of the protein derivative in an alkaline buffer which further contains a zinc salt,
- b) adjusting the pH value of the solution to a value between 7 and 10 and
- c) isolating the crystals formed.

5

In a preferred embodiment, the protein derivative which has a lysine residue which carries a lipophilic substituent on the  $\epsilon$ -amino group is a proinsulin, an insulin or an insulin analogue in which the  $\epsilon$ -amino group of a lysine residue carries an acyl group.

- 10 In another preferred embodiment, the protein derivative is an insulin derivative selected from the group comprising N<sup>B29</sup>-(myristoyl)des(B30) human insulin, N<sup>B29</sup>-(myristoyl) human insulin, N<sup>B29</sup>-(palmitoyl) human insulin, N<sup>B29</sup>-(myristoyl)Lys<sup>B29</sup>Pro<sup>B29</sup> human insulin, N<sup>B29</sup>-(palmitoyl)Lys<sup>B29</sup>Pro<sup>B29</sup> human insulin, N<sup>B30</sup>-(myristoyl)Thr<sup>B29</sup>Lys<sup>B30</sup> human insulin, N<sup>B30</sup>-(palmitoyl)Thr<sup>B29</sup>Lys<sup>B30</sup> human insulin, N<sup>B29</sup>-(N-palmitoyl- $\gamma$ -glutamyl)des(B30) human insulin,
- 15 N<sup>B29</sup>-(N-lithocholyl- $\gamma$ -glutamyl)des(B30) human insulin and N<sup>B29</sup>-( $\omega$ -carboxyheptadecanoyl)des(B30) human insulin.

In another preferred embodiment, a phenol is added to the solution before the final adjustment of the pH value.

20

In another preferred embodiment, an amount of phenol added which is from about 1.5% (w/w) to about 10% (w/w), preferably from about 1.5% (w/w) to about 3% (w/w) of the amount of protein present in the solution.

- 25 In another preferred embodiment, a phenol selected from the group comprising hydroxybenzene, *m*-cresol, methylparabene and ethylparabene is added.

In another preferred embodiment, the lipophilic group on the  $\epsilon$ -amino group of the lysine residue is an acyl group having from 4 to 40, more preferred from 10 to 40 carbon atoms.

30

In another preferred embodiment, the lipophilic group on the  $\epsilon$ -amino group of the lysine residue is a straight chain acyl group.

In another preferred embodiment, the buffer is composed of ammonia or an amine and an acid.

In another preferred embodiment, the buffer is composed of ammonia and phosphoric acid.

5

In another preferred embodiment, the buffer is composed of ammonia and a carboxylic acid.

In another preferred embodiment, the buffer is composed of an amine and phosphoric acid.

10 In another preferred embodiment, the buffer is composed of an amine and a carboxylic acid.

In another preferred embodiment, when the buffer is composed of an amine and an acid, the amine is selected from the group comprising tris(hydroxymethyl)aminomethane, 2-amino-2-methyl-1,3-propanediol, 2-hydroxyethylamine and tris(2-hydroxyethyl)amine.

15

In another preferred embodiment, the buffer is composed of an ampholytic compound, preferably aspartic acid or N-tris(hydroxymethyl)methylglycine, and optionally an acid.

20 In another preferred embodiment, when the buffer is composed of ammonia and a carboxylic acid or an amine and a carboxylic acid, the carboxylic acid is selected from the group comprising acetic acid, citric acid, lactic acid, malic acid, malonic acid, succinic acid, tartaric acid, tartronic acid and tricarballic acid.

25 In another preferred embodiment, the concentration of the ammonia or amine in the buffer is between 0.1 M and 1 M, preferably between 0.2 M and 0.6 M.

In another preferred embodiment, the concentration of the phosphoric acid or the carboxylic acid in the buffer is between 0.05 M and 0.5 M, preferably between 0.05 M and 0.2 M.

30 In another preferred embodiment, the zinc salt added to the solution is zinc chloride or a zinc salt of a carboxylic acid, preferably a zinc salt of one the following acids: acetic acid, citric acid, lactic acid, malic acid, malonic acid, succinic acid, tartaric acid, tartronic acid and tricarballic acid.

In another preferred embodiment, the zinc salt is added in a molar amount which is between 33% and 150% of the molar amount of the peptide monomer.

5 In another preferred embodiment, wherein the protein derivative which has a lysine residue which carries a lipophilic substituent on the  $\epsilon$ -amino group is a proinsulin, an insulin or an insulin analogue in which the  $\epsilon$ -amino group of a lysine residue carries an acyl group, the pH value of the solution in which the precipitation of crystals is performed is adjusted to a value in the range from about 8.0 to about 8.5.

10

## DETAILED DESCRIPTION OF THE INVENTION

### Definitions

As used in the present text the designation "insulin" is used to designate any naturally occurring insulin. The designation "insulin analogue" is used to designate a peptide with insulin activity, formally derived from a naturally occurring insulin by exchange of one or more amino acid residues and/or deletion of one or more amino acid residues and/or addition of one or more amino acid residues. An "acylated insulin" (or insulin analogue) is an insulin (or insulin analogue) which has an acyl group in the  $\epsilon$ -amino group of a lysine residue contained in said insulin (or insulin analogue).

20

### Preferred embodiments

The precipitation of the crystals according to the present method is carried out in water which optionally contains a co-solvent. When a co-solvent is used, this is preferably a water-miscible solvent e.g. an alcohol.

25

The operations preceding the precipitation of the crystals are preferably carried out at a temperature around ambient. After standing from about 2 hours to about 40 hours at about ambient, the reaction mixture is cooled to a temperature near 0° C until no further precipitation of crystals occurs and the crystals are collected on a filter. If desired, the crystals can be washed with an ice-cold buffer solution corresponding to the solution from which the precipitation took place and with ice-cold ethanol.

30

The components of the buffer solution should preferably be less prone to form zinc complexes than insulin.

The present invention is further illustrated by the following examples which, however, are not to be construed as limiting the scope of protection as described in the appended claims. The features disclosed in the foregoing description and in the following examples may, in any combination thereof, be material for realizing the invention in diverse forms thereof.

10

## EXAMPLES

### Example 1

Preparation of crystals of N<sup>B29</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

15

Three stock solutions a), b) and c) were used during the preparation of crystals of N<sup>B29</sup>-(myristoyl)des(B30) human insulin. These solutions were prepared as follows:

#### Stock solution a)

20

12.11 g of tris(hydroxymethyl)aminomethane was dissolved in 80 ml of water and the pH value of the solution was adjusted to 8.30 by means of approximately 7 ml of 5 N hydrochloric acid. Water was then added to a final volume of 100 ml.

#### Stock solution b)

25

14.71 g of trisodium citrate dihydrate and 0.11 g of zinc acetate dihydrate were dissolved in water and 6.25 ml of a 3% (w/v) solution of phenol in water was added and the final volume of the solution adjusted to 100 ml.

#### Stock solution c)

30

1.471 g of trisodium citrate dihydrate and 2.422 g of tris(hydroxymethyl)aminomethane were dissolved in water and the volume of the solution adjusted to 40 ml. The pH value was adjusted to 8.1 using 5 N hydrochloric acid and finally the volume was adjusted to 50 ml with water.



Crystallization of N<sup>1829</sup>-(myristoyl)des(B30) human insulin

1.00 g of amorphous N<sup>1829</sup>-(myristoyl)des(B30) human insulin powder, obtained as described in WO 95/07931, was dispersed in a mixture of 38 ml of water and 2 ml of absolute ethanol and 40 ml of stock solution a) was added with stirring. When the insulin had dissolved 20 ml of stock solution b) was added and the pH value of the mixture was adjusted to be in the range 8.1-8.2 by addition of approximately 1 ml of 5 N hydrochloric acid. The mixture was left at ambient temperature with slow stirring overnight and then cooled to 4° C. The crystals formed were collected on a 50 mm filter and quickly washed with ice-cold stock solution c). After draining, the crystals were washed with 20 ml of ice-cold absolute ethanol and the drained crystals were dried *in vacuo*.

**Example 2**

Preparation of crystals of N<sup>1829</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of 2-amino-2 methyl-1,3-propanediol in place of tris(hydroxymethyl)aminomethane.

**Example 3**

Preparation of crystals of N<sup>1829</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of 2-hydroxyethylamine in place of tris(hydroxymethyl)aminomethane.

**Example 4**

Preparation of crystals of N<sup>1829</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of 2-amino-2 methyl-1,3-propanediol in place of tris(hydroxymethyl)aminomethane.

**Example 5**

Preparation of crystals of N<sup>1829</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of ammonia in place of tris(hydroxymethyl)aminomethane.

**Example 6**

Preparation of crystals of N<sup>629</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of tris(2-hydroxyethyl)amine in place of tris(hydroxymethyl)aminomethane.

**Example 7**

Preparation of crystals of N<sup>629</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of N-tris(hydroxymethyl)methyl-glycine in place of tris(hydroxymethyl)aminomethane.

**Example 8**

Preparation of crystals of N<sup>629</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of L-aspartic acid in place of tris(hydroxymethyl)aminomethane.

**Example 9**

Preparation of crystals of N<sup>629</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of L-aspartic acid in place of both tris(hydroxymethyl)aminomethane and citric acid.

**Example 10**

Preparation of crystals of N<sup>629</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of sodium acetate in place of trisodium citrate.

**Example 11**

Preparation of crystals of N<sup>629</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of disodium tartrate in place of trisodium citrate.

**Example 12**

Preparation of crystals of N<sup>B29</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of disodium succinate in place of trisodium citrate.

**Example 13**

Preparation of crystals of N<sup>B29</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of disodium hydrogen phosphate in place of trisodium citrate.

**Example 14**

Preparation of crystals of N<sup>B29</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of disodium malate in place of trisodium citrate.

**Example 15**

Preparation of crystals of N<sup>B29</sup>-(myristoyl)des(B30) human insulin containing zinc and phenol.

The crystallization procedure according to Example 1 was repeated with use of disodium malonate in place of trisodium citrate.

**Example 16**

Crystallization of N<sup>B29</sup>-(myristoyl)Lys<sup>B28</sup>Pro<sup>B29</sup> human insulin.

The crystallization procedure according to Example 1 was repeated with use of N<sup>B29</sup>-(myristoyl)Lys<sup>B28</sup>Pro<sup>B29</sup> human insulin in place of N<sup>B29</sup>-(myristoyl)des(B30) human insulin.

**Example 17**

Crystallization of N<sup>B29</sup>-( $\omega$ -carboxyheptadecanoyl)des(B30) human insulin.

The crystallization procedure according to Example 1 was repeated with use of N<sup>B29</sup>-( $\omega$ -carboxyheptadecanoyl)des(B30) human insulin in place of N<sup>B29</sup>-(myristoyl)des(B30) human insulin.

**Example 18**

Crystallization of N<sup>B29</sup>-(N-lithocholyl- $\gamma$ -glutamyl)des(B30) human insulin.

The crystallization procedure according to Example 1 was repeated with use of N<sup>B29</sup>-(N-lithocholyl-γ-glutamyl)des(B30) human insulin in place of N<sup>B29</sup>-(myristoyl)des(B30) human insulin.

5 **Example 19**

**Crystallization of N<sup>B29</sup>-(myristoyl)des(B30) human insulin**

The crystallization method according to Example 1 was performed with use of sodium chloride in place of tris(hydroxymethyl)aminomethane following the procedure described below.

- 10 Two stock solutions d) and e) were used during the crystallization procedure. These solutions were prepared as follows:

**Stock solution d)**

- 15 14.71 g of trisodium citrate dihydrate, 5.844 g of sodium chloride and 0.11 g of zinc acetate dihydrate were dissolved in about 75 ml of water and after addition of 6.25 ml of a 3% (w/v) solution of phenol in water the pH value was adjusted to 8.1 using 2-N sodium hydroxide and the final volume of the solution was adjusted to 100 ml.

**Stock solution e)**

- 20 1.471 g of trisodium citrate dihydrate and 2.922 g of sodium chloride were dissolved in about 35 ml of water. The pH value was adjusted to 8.1 using 2 N sodium hydroxide and finally the volume was adjusted to 50 ml with water.

**Crystallization procedure:**

- 25 1.00 g of amorphous N<sup>B29</sup>-(myristoyl)des(B30) human insulin powder was dispersed in a mixture of 78 ml of water and 2 ml of absolute ethanol and pH was adjusted to 8.3 with 0.1 M NaOH. When the insulin had dissolved 20 ml of a stock solution d) was added and the pH value of the mixture was adjusted to be in the range 8.1-8.2 by addition of 1 N hydrochloric acid. The mixture was slowly stirred overnight at ambient temperature and then cooled to 4°
- 30 C. The crystals formed were collected on a 50 mm planar filter and quickly washed with ice-cold stock solution e). After draining, the crystals were washed with 20 ml of ice-cold absolute ethanol and the drained crystals were dried *in vacuo*.

## CLAIMS

1. A method of providing zinc containing crystals of a protein derivative which has a lysine residue which carries a lipophilic substituent on the  $\epsilon$ -amino group, said method comprising
  - a) providing a solution of the protein derivative in an alkaline buffer, which further contains a zinc salt,
  - b) adjusting the pH value of the solution to a value between 7 and 10 and
  - c) isolating the crystals formed.
2. The method of claim 1 wherein the protein derivative is proinsulin, insulin or an insulin analogue in which the  $\epsilon$ -amino group of a lysine residue carries an acyl group.
3. The method of claim 1 wherein the protein derivative is an insulin derivative selected from the group comprising N<sup>B29</sup>-(myristoyl)des(B30) human insulin, N<sup>B29</sup>-(myristoyl) human insulin, N<sup>B29</sup>-(palmitoyl) human insulin, N<sup>B29</sup>-(myristoyl)Lys<sup>B29</sup>Pro<sup>B29</sup> human insulin, N<sup>B29</sup>-(palmitoyl)Lys<sup>B29</sup>Pro<sup>B29</sup> human insulin, N<sup>B30</sup>-(myristoyl)Thr<sup>B29</sup>Lys<sup>B30</sup> human insulin, N<sup>B30</sup>-(palmitoyl)Thr<sup>B29</sup>Lys<sup>B30</sup> human insulin, N<sup>B29</sup>-(N-palmitoyl- $\gamma$ -glutamyl)-des(B30) human insulin, N<sup>B29</sup>-(N-lithocholyl- $\gamma$ -glutamyl)des(B30) human insulin and N<sup>B29</sup>-( $\omega$ -carboxyheptadecanoyl)des(B30) human insulin.
4. The method of claim 1 wherein a phenol is added to the solution before the final adjustment of the pH value.
5. The method of claim 4 wherein the amount of phenol added is from about 1.5% (w/w) to about 10% (w/w), preferably from about 1.5% (w/w) to about 3% (w/w) of the amount of protein present in the solution.
6. The method of claim 4 wherein the phenol is selected from the group comprising hydroxybenzene, *m*-cresol, methylparabene and ethylparabene.
7. The method of claim 1 wherein the lipophilic group is an acyl group having from 4 to 40, more preferred from 10 to 40 carbon atoms.
8. The method of claim 1 wherein the lipophilic group is a straight chain acyl group.
9. The method of claim 1 wherein the alkaline buffer is a nitrogen-containing buffer.
10. The method of claim 9 wherein the buffer is composed of ammonia or an amine and an acid.
11. The method of claim 9 wherein the buffer is composed of ammonia and phosphoric acid.

12. The method of claim 9 wherein the buffer is composed of ammonia and a carboxylic acid.
13. The method of claim 9 wherein the buffer is composed of an amine and phosphoric acid.
- 5 14. The method of claim 9 wherein the buffer is composed of an amine and a carboxylic acid.
15. The method of claim 9 wherein the buffer is composed of an ampholytic compound, preferably aspartic acid or N-tris(hydroxymethyl)methylglycine, and optionally an acid.
16. The method of claim 13 or 14 wherein the amine is selected from the group comprising  
10 tris(hydroxymethyl)aminomethane, 2-amino-2-methyl-1,3-propanediol, 2-hydroxyethylamine and tris(2-hydroxyethyl)amine.
17. The method of claim 12 or 14 wherein the carboxylic acid is selected from the group comprising acetic acid, citric acid, lactic acid, malic acid, malonic acid, succinic acid, tartaric acid, tartronic acid and tricarballic acid.
- 15 18. The method of claim 10 wherein the concentration of the ammonia or amine in the buffer is between 0.1 M and 1 M, preferably between 0.2 M and 0.6 M.
19. The method of claim 10 wherein the concentration of the phosphoric acid or the carboxylic acid in the buffer is between 0.05 M and 0.5 M, preferably between 0.05 M and 0.2 M.
- 20 20. The method of claim 1 wherein the zinc salt added to the solution is zinc chloride or a zinc salt of a carboxylic acid, preferably a zinc salt of one the acids mentioned in claim 17.
21. The method of claim 1 wherein the zinc salt is added in a molar amount which is between 33% and 150% of the molar amount of the peptide monomer.
- 25 22. The method of claim 1 wherein the pH value is adjusted to a value in the range from about 8.0 to about 8.5.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 98/00042

<b>A. CLASSIFICATION OF SUBJECT MATTER</b>		
IPC6: C07K 14/62, C07K 1/30 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b>		
Minimum documentation searched (classification system followed by classification symbols)		
IPC6: C07K		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
SE,DK,FI,NO classes as above		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
REG, CAPLUS, WPI		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0747391 A2 (ELI LILLY AND COMPANY), 11 December 1996 (11.12.96)	1-22
Y	US 4959351 A (ULRICH GRAU), 25 Sept 1990 (25.09.90), see claims	1-22
A	US 3856771 A (RICHARD L. JACKSON), 24 December 1974 (24.12.74)	1-22
A	GB 2290294 A (ELI LILLY AND COMPANY), 20 December 1995 (20.12.95)	1-22
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "A" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
6 May 1998		20.05.98
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86		Authorized officer Carolina Gómez Lag r18f Telephone No. +46 8 782 25 00

Form PCT/ISA/210 (second sheet) (July 1992)

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

02/04/98

International application No.

PCT/DK 98/00042

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0747391 A2	11/12/96	AU 5951396 A	30/12/96
		NO 975583 A	03/12/97
		US 5700904 A	23/12/97
		WO 9640730 A	19/12/96
US 4959351 A	25/09/90	AU 567360 B	19/11/87
		AU 3127284 A	31/01/85
		CA 1246549 A	13/12/88
		DE 3327709 A	07/02/85
		DE 3469535 A	07/04/88
		DK 172241 B	02/02/98
		DK 368884 A	30/01/85
		EP 0133285 A,B	20/02/85
		SE 0133285 T3	
		FI 842981 A	30/01/85
		JP 2029981 C	19/03/96
		JP 7064876 B	12/07/95
		JP 60051118 A	22/03/85

Form PCT/ISA/210 (patent family annex) (July 1992)



**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

02/04/98

International application No.  
PCT/DK 98/00042

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 3856771 A	24/12/74	AR 199541 A	09/09/74
		AT 344925 B	25/08/78
		AU 6959774 A	04/12/75
		BE 518137 A	00/00/00
		BE 522895 A	00/00/00
		BE 817649 A	15/01/75
		BG 25803 A	12/12/78
		CA 1025851 A	07/02/78
		CH 197889 A	00/00/00
		CH 313787 A	00/00/00
		CH 319968 A	00/00/00
		CH 620362 A	28/11/80
		CS 183753 B	31/07/78
		DD 114597 A	12/08/75
		DE 728213 C	00/00/00
		DE 730948 C	00/00/00
		DE 1009057 B	00/00/00
		DE 1118651 B	00/00/00
		DE 2431483 A,B,C	06/02/75
		DK 148282 B,C	28/05/85
		DK 379374 A	17/03/75
		FR 816799 A	17/08/37
		FR 1076287 A	25/10/54
		FR 2237879 A,B	14/02/75
		GB 487534 A	00/00/00
		GB 747968 A	00/00/00
		GB 748007 A	00/00/00
		GB 890830 A	00/00/00
		GB 1472427 A	04/05/77
		JP 1148564 C	26/05/83
		JP 50040717 A	14/04/75
		JP 57035825 B	31/07/82
		NL 56198 C	00/00/00
		NL 57064 C	00/00/00
		NL 67951 C	00/00/00
		NL 82149 C	00/00/00
		NL 92858 C	00/00/00
		NL 102298 C	15/03/62
		NL 179910 B,C	01/07/86
		NL 253505 A	00/00/00
		NL 7409501 A	20/01/75
		SE 415229 B,C	22/09/80
		SE 7409248 A	17/01/75
		US 2762337 A	11/09/56
		ZA 7403424 A	28/01/76

Form PCT/ISA/210 (patent family annex) (July 1992)

**INTERNATIONAL SEARCH REPORT**  
Information on patent family members

02/04/98

International application No.  
PCT/DK 98/00042

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB 2290294 A	20/12/95	AU 2168295 A	04/01/96
		BE 1009409 A	04/03/97
		BR 9502797 A	12/03/96
		CA 2151564 A	17/12/95
		CN 1116629 A	14/02/96
		CZ 9501543 A	14/02/96
		DE 19521753 A	21/12/95
		DK 67695 A	17/12/95
		ES 2091728 A	01/11/96
		FI 952932 A	17/12/95
		FR 2721215 A,B	22/12/95
		GB 9512105 D	00/00/00
		HU 71908 A	28/02/96
		HU 9501717 D	00/00/00
		IE 68852 B	24/07/96
		IE 950435 A	27/12/95
		IL 114153 D	00/00/00
		IT 1276722 B	03/11/97
		IT M1951277 A	18/12/95
		JP 8003064 A	09/01/96
		LU 88627 A	01/02/96
		NL 1000565 A,C	00/00/00
		NL 1004643 A,C	00/00/00
		NO 952356 A	18/12/95
		NZ 272360 A	24/02/97
		PL 309100 A	27/12/95
		PT 101723 A,B	29/12/95
		SE 9502168 A	17/12/95
		SI 9500199 A	29/02/96
		US 5461031 A	24/10/95
		US 5650486 A	22/07/97

Form PCT/ISA/210 (patent family annex) (July 1992)